

Translator: Lucia L. Severinghaus
Date: September 8, 1981

Acta Zootaxonomica Sinica 6(1): 97

A NEW RECORD OF CRANE FROM CHINA

In January 1979, the Kun-ming City Zoo (Yun-nan Province) was collecting specimens in Shu County, Chiang-su Province. The collecting team purchased a crane which is identified to be Grus canadensis Linnaeus. This is a new record for China.

Measurements (mm): Live bird, length 850, wing 457, tail 190, tarsus (tarsometatarsus) 180, midtoe (including nail) 81.

Descriptions: Upper body slate grey. Wing coverts lighter grey with rusty brown borders. Feather shafts black. Tail feather slaty grey. Neck entirely greyish blue. White below the eyes, on cheek, and throat. Breast and belly light brown with grey borders on the feathers. Lores and the naked part on the head vermillian red. The border of the naked and the feathered portions on the head is shaped like a blunt "v". Beak black, upper mandible lighter colored. Feet and toes black. Nictitating membrane deep red.

Distribution: Siberia, Alaska, south to Mexico. In China, found in Chiang-su Province, probably a stray bird.

Kuang Bang-yü (China National Academy
of Science, Institute of Zoology)
Hsian Ju-lun (Kun-ming City Zoo)
Wang Tse-chiang (Yun-nan University,
Biology Department)

布的雾点是暗红褐色,而不是灰褐色;雄性外生殖器的抱器瓣只有中央一个小突起,阳茎上的齿形突起不在中央而在末端之前。

A NEW SPECIES OF *SUZUKIA* MATSUMURA (LEPIDOPTERA: NOTODONTIDAE)

CAI RONG-QUAN

(Institute of Zoology, Academia Sinica)

Suzukia sichuanensis, sp. nov.

Expanse of ♂ forewing 43—46 mm. Very similar in appearance to *S. cinerea* (Butler), but distinguishable from it by the following characters: ground-colour of forewing white, with dark reddish-brown dusts; valve with a single median process; aedeagus with a dentiform subterminal process.

Holotype ♂, Emei Shan, Sichuan, 800—1000 m, 1.V.1957.

Paratypes 2♂♂, ditto; 1♂, Xichang (Lu Shan), Sichuan, 1800 m, 3.VIII.1974.

中 国 鹤 类 新 纪 录

A NEW RECORD OF CRANE FROM CHINA

云南省昆明市动物园 1979 年 1 月在江苏省沭阳县征集展出动物时,购得鹤类一只。经鉴定为 *Grus canadensis* Linnaeus, 系我国鹤科鸟类种的新纪录。现提出报道。

量度(毫米): 1♀(活鸟),全长 850,翅长 457,尾长 190,附蹼 180,中趾(连爪) 81。

形态: 上体几为石板灰色。翼上覆羽较淡,羽缘锈棕色,羽轴深黑色;尾羽石板灰色;整个颈部都为灰蓝色;眼下、颊、颏及喉部为白色;胸部及腹部浅棕色,且具灰色羽缘。眼先及头部裸出部分为朱红色;裸出部与头顶的纤羽区的交界处呈钝角“V”形。嘴黑色,上喙较浅。脚和趾黑色。虹膜深红色。

分布: 在国外分布于西伯利亚,阿拉斯加,直到墨西哥。在我国江苏省的发现,应为迷鸟。

匡邦郁

鲜汝伦

王紫江

(中国科学院昆明动物研究所) (昆明市动物园) (云南大学生物系)